




ASTRONOMY OPEN NIGHT | 23RD MAY 2015



CAN BLACK HOLES SHINE?

MICHAEL COWLEY

TIMELINE OF BLACK HOLES


- 1915: Einstein's Theory of General Relativity
- 1919: Eddington's solar eclipse experiment
- 1967: Wheeler coins the term "black hole"
- 1970s: Hawking advances our understanding of black holes
- 2002: The Milky Way's lurking monster
- Today: What do we know and how do we find them?







1915: Albert Einstein's General Relativity

Albert Einstein's general theory of relativity describes the interaction of gravity as a result of space being curved by massive objects



1919: Sir Arthur Eddington's Solar Eclipse Experiment

Eddington photographed positions of stars near the Sun to test Albert Einstein's prediction of the bending of light around massive objects from his general theory of relativity




EDDINGTON'S ACTUAL ECLIPSE PHOTO
MAY 29, 1919

An unimaginably dense region of space where space is curved around it so completely and gravity becomes so strong that nothing, not even light, can escape.




1964: John Wheeler coins the term “black hole”

American theoretical physicist, John Wheeler helps popularise the study of general relativity in the mainstream of theoretical physics, and coins the term “black holes”




1970s: Stephen Hawking Advances our Understanding of Black Holes

In 1968, Stephen Hawking joins the Institute of Astronomy in Cambridge and begins to apply the laws of thermodynamics and quantum mechanics to black holes




BIRTH OF A BLACK HOLE




The fate of a star depends on its mass (size not to scale)

BLACK HOLE TYPES

- **Stellar-mass**: black holes with 3 to 20 times the mass of our own Sun
- **Supermassive**: black holes with millions to billions of times the mass of our own Sun





2002: The Milky Way's
Lurking Monster

Black Holes

How do we find them?



3C 273

QUASAR 3C 273




4 TRILLION
TIMES BRIGHTER
THAN THE SUN!

- Discovered in 1959
- **273rd** object in the **3rd C**ambridge Radio Survey (**3C 273**)
- Classified as a Quasar or quasi-stellar radio source (looks like a star, but is not)
- Approximately two billion light years away (that's 2×10^{22} or 20,000,000,000,000,000,000 kilometres away)




How Bright is it?

ELECTROMAGNETIC SPECTRUM




QUASAR 3C 273 SPECTRUM



Black Holes


Where does the light come from?



- Gravity causes material to spiral inward towards the black hole
- Frictional forces compress and raise the temperature of the material causing the emission of light ranging from X-rays to infrared
- Particles accelerated to speeds approaching that of light and emerge from the poles as radio jets

QUASAR 3C 273 RADIO JET

~2 x longer than
distance across
Milky Way



CAN BLACK HOLES SHINE?



**4 TRILLION
TIMES BRIGHTER
THAN THE SUN!**



THANK YOU!

Want to learn more?

QR CODE
SCAN ME!

<http://hubblesite.org>



HUBBLESITE Special Feature

BLACK HOLES: Gravity's Relentless Pull

Home **Journey to a Black Hole** **Black Hole Encyclopedia**

Finding the Invisible **The Voyage** **Up Close and Personal**

Speed **Distance from Earth**

99.999...95%
99.999...95%
99.999...95%
99.9995%
90%
lightspeed

1 million
10,000
1000
1 Light Years

Replay last part of voyage

You have reached escape velocity, but at this speed it will take you days to reach the moon ... and over 16 years to reach Pluto. Please use your engine to generate more energy for your spacecraft.

1 billion times more energy 10 billion times more energy 100 billion times more energy 1,000 billion times more energy

Content Overview **Credits** **Contact Us** **Glossary** **How to Print**

